

Marshall University
Marshall Digital Scholar

Theses, Dissertations and Capstones

2015

Effects of Roads and Trails on Peaks of Otter Salamander (*Plethodon hubrichti*) and Eastern Red-backed Salamander (*Plethodon cinereus*) Movement Behavior

Cory Benjamin Goff
goff57@marshall.edu

Follow this and additional works at: <http://mds.marshall.edu/etd>



Part of the [Animal Sciences Commons](#), and the [Behavior and Ethology Commons](#)

Recommended Citation

Goff, Cory Benjamin, "Effects of Roads and Trails on Peaks of Otter Salamander (*Plethodon hubrichti*) and Eastern Red-backed Salamander (*Plethodon cinereus*) Movement Behavior" (2015). *Theses, Dissertations and Capstones*. Paper 951.

This Thesis is brought to you for free and open access by Marshall Digital Scholar. It has been accepted for inclusion in Theses, Dissertations and Capstones by an authorized administrator of Marshall Digital Scholar. For more information, please contact zhangj@marshall.edu.

EFFECTS OF ROADS AND TRAILS ON PEAKS OF OTTER SALAMANDER
(*PLETHODON HUBRICHTI*) AND EASTERN RED-BACKED SALAMANDER
(*PLETHODON CINEREUS*) MOVEMENT BEHAVIOR

A thesis submitted to
the Graduate College of
Marshall University
In partial fulfillment of
the requirements for the degree of
Master of Science
in
Biological Sciences: Organismal, Evolutionary and Ecological Biology
by
Cory Benjamin Goff
Approved by
Dr. Jayme L. Waldron, Committee Chairperson
Dr. Anne C. Axel
Dr. Thomas K. Pauley
Dr. Shane M. Welch

Marshall University
August 2015

©2015
Cory Benjamin Goff
ALL RIGHTS RESERVED

ACKNOWLEDGMENTS

First, I would like to thank God, my Savior and Creator, for the desire to learn about the natural world around me and to be a good steward of the precious wildlands and wildlife that I have come to appreciate and enjoy. Second, I would like to thank my undergraduate advisor, Dr. Norm Reichenbach at Liberty University, for giving me the instruction and experience with herpetofauna that led to my love for reptiles and amphibians. Also, his guidance led me to continue my education, pursue a Master's degree, and stick with research in the field of herpetology. I would also like to thank my advisor here at Marshall, Dr. Jayme Waldron, for accepting me into her herpetology lab and for her continued support and guidance in my research. Her knowledge of statistics and desire for excellence motivated me to learn valuable techniques, gain necessary experience, and press on in my research. Her continued assistance in writing style and word usage has made me a better writer. I would like to thank my committee members, Dr. Anne Axel, Dr. Thomas Pauley, and Dr. Shane Welch for their continued help and guidance. Dr. Axel was very valuable in aspects of image processing and data collection. Her experience in the programs I used was key to help me measure distances from digital imagery. Dr. Pauley was always a source of encouragement and an encyclopedia of knowledge. His understanding of salamander life history and memory of reference papers was exceptional. Dr. Welch always had a new way to approach the project and widen my focus. His broad knowledge of research and study designs helped me construct my own project. In addition, his statistical knowledge combined with that of everyone else helped me examine new angles.

I would like to thank the Herpetology Lab members, both past and present, for their encouragement and support through my time at Marshall. Derek Breakfield, Elise Edwards and Kelli Herrick, as second year students when I arrived, thank you for the wise counsel, all the fun

times together, and for just looking out for us “kids” in the lab. Jonathan Cooley, Bradley O’Hanlon, and Shelby Timm, you were all an encouragement to me and as we struggled through graduate life together, it was great to have friends there all the way through. The two years together went by so fast, but at the same time, I feel I have known you all for so long and I am very thankful for the friendships made. Jessyca Conatser, Matt Grisnik, Berlynn Heres, and Theresa Houze, as our “progeny” I hope we have helped you and prepared you well. Good luck with all your studies and research and thank you for the fun times and learning together. I would also like to thank Nathan Fleshman for his significant role in helping me with my field data collection over the summer, and working with me to get my research up and running. It was a huge encouragement to have someone with me the first few weeks of being out in the field in the wee hours of the night.

I also want to thank my wife, Beth, and my good friend and roommate, Thaddaeus Tuggle. Beth has been a huge supporter of my work and research and has always been there for me, encouraging me to press on and stay focused. I thank her for her love toward me and her desire to see me succeed. She has joined me on several “herping” adventures and has always been a faithful companion. As one of the first people I met when I arrived, Thaddaeus has been an encouragement in the science department and has become a great friend. He has also accompanied me on several adventures during my time here and always had a way to make things fun and entertaining.

Finally, I would like to thank my parents, for raising me to appreciate and be curious about the natural world, and to work hard to achieve my goals. They have been a constant support and without their love and help, I would not be where I am today. They have sacrificed much to make my life better and to see me succeed.

TABLE OF CONTENTS

Acknowledgements.....	iii
Table of Contents.....	v
List of Tables.....	vii
List of Figures.....	x
Abstract.....	xi
Chapter I: Effects of Roads and Trails on Peaks of Otter Salamander (<i>Plethodon hubrichti</i>) and Eastern Red-backed Salamander (<i>Plethodon cinereus</i>) Movement Behavior.....	1
Introduction.....	1
Methods.....	4
Study Sites.....	4
Data Collection.....	5
Data Analysis.....	7
Candidate Models.....	8
Statistical Analysis.....	9
Results.....	10
Correlation Analysis.....	12
Movement Probability.....	13
Peaks of Otter Salamander.....	13
Red-backed Salamander.....	15
Total Distance Moved.....	16
Peaks of Otter Salamander.....	16

Red-backed Salamander.....	18
Net Distance Moved.....	19
Peaks of Otter Salamander.....	19
Red-backed Salamander.....	20
Movement Direction.....	21
Peaks of Otter Salamander.....	21
Red-backed Salamander.....	22
Discussion.....	23
Literature Cited.....	28
Appendix A.....	32
Salamander Crossing Note.....	32
Salamander Microhabitat Use.....	32
IACUC Approval Letter- Peaks of Otter Salamander.....	34
IACUC Approval Letter- Eastern Red-backed Salamander.....	35

LIST OF TABLES

1	Definition of predictor variables used in Peaks of Otter Salamander and Eastern Red-backed Salamander movement behavior models.....	9
2	Average mass (g) and snout-vent length (SVL; mm) for Peaks of Otter salamander (POS) and Eastern Red-backed salamander (RBS) with standard deviation and range of values.....	11
3	Average total distance moved (m) and net distance moved (m) for Peaks of Otter salamanders (POS) and Eastern Red-backed salamanders (RBS), including range and standard deviation.....	12
4	Predictor variables used in candidate models to analyze Peaks of Otter Salamander and Eastern Red-backed Salamander movement behavior.....	13
5	Candidate models examining Peaks of Otter Salamander (N = 141) movement probability (binary logistic regression), ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....	14
6	Candidate models examining Eastern Red-backed Salamander (N = 46) movement probability (binary logistic regression), ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....	16
7	Candidate models examining Peaks of Otter Salamander (N = 141) total distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current	

- model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....17
- 8 Candidate models examining Eastern Red-backed Salamander (N = 46) total distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....18
- 9 Candidate models examining Peaks of Otter Salamander (N = 141) net distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....19
- 10 Candidate models examining Eastern Red-backed Salamander (N = 46) net distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....20
- 11 Candidate models examining Peaks of Otter Salamander (N = 141) movement direction (multinomial logistic regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....21

12	Candidate models examining Eastern Red-backed Salamander (N = 46) movement direction (multinomial logistic regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. $\Delta AICc$ = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta AICc < 2.00$ (in bold) were retained for inference.....	22
----	---	----

LIST OF FIGURES

1	Fluorescent pigment powder applied to the belly and tail of a salamander as it looks in natural light (A) and under ultraviolet light (B).....	6
2	Georectified image of a marked salamander movement trail using reflective rope with 1m ² PVC grid, north arrow (orange), and dry-erase board with identification information and direction of treatment edge (blue arrow).....	7
3	Diagram of movement direction split into three equal bearings for analysis.....	8
4	Average distance moved (meters) for each species with pooled data for gender and treatment.....	12
5	A Peaks of Otter Salamander perched on a small plant, covered in fluorescent pigment powder used for tracking.....	33
6	A Peaks of Otter Salamander climbing the trunk of a tree.....	33

ABSTRACT

Habitat fragmentation negatively affects movement and dispersal of wildlife by altering useable space and creating barriers. The objective of this study was to assess impacts of fragmentation by roads and trails on the movement behavior of two terrestrial, woodland salamander species. I evaluated whether proximity to roads and trails affected the movement behavior of two species: the Peaks of Otter Salamander (*Plethodon hubrichti*), a species of special concern, and the Eastern Red-backed Salamander (*Plethodon cinereus*). I examined movement behavior in three treatments: paved roads, gravel roads and hiking trails. I compared treatments to controls located at least 200m from treatment edges to assess movement behavior based on fragmentation type. I detected salamanders by flipping cover objects during the day. I tracked movement patterns using a fluorescent pigment powder applied to each individual. I mapped and measured each powder trail to quantify travel distance and direction. I analyzed treatment and abiotic effects on movement behavior. I failed to detect a significant effect of treatment or distance from the treatment edge on individual foraging movements. Analysis indicated that abiotic factors including moon phase and precipitation were significant predictors of movement. This study provided additional insight into plethodontid movement patterns. Research will require managers to consider road and trail effects when designing and implementing long-term management plans for the survival of species within the Peaks of Otter area and other regions of the Appalachian Mountains.

CHAPTER 1

EFFECTS OF ROADS AND TRAILS ON PEAKS OF OTTER SALAMANDER (*PLETHODON HUBRICHTI*) AND EASTERN RED-BACKED SALAMANDER (*PLETHODON CINEREUS*) MOVEMENT BEHAVIOR

INTRODUCTION

Fragmentation, the process of breaking contiguous habitat into smaller sections, is a problem when managing wildlife populations as new roads are built and habitats are altered for forest management (Forman and Alexander 1998). Roads create a barrier effect, fragmenting continuous populations into smaller, partially isolated subpopulations (Forman and Alexander 1998). Such subpopulations may decrease along with patch size as habitat quality declines (Semlitsch et al. 2007). Habitat alterations and perceived risk associated with roads and other barriers affect movement and dispersal of various taxa (Mazerolle 2001; Marsh and Beckman 2004; Marsh et al. 2005; Semlitsch et al. 2007; Connette and Semlitsch 2013). In addition, roads and trails facilitate anthropogenic activity including vehicular and foot traffic, which also acts as barriers to animal movements (Andrews et al. 2008; Balkenhol and Waits 2009).

Habitat patches and decreased habitat suitability created by edges limit dispersal and movement for many fauna (Marsh et al. 2005; Semlitsch et al. 2007; Row et al. 2012). Roads cause direct and indirect mortality of terrestrial animals, affecting population demographics (Andrews et al. 2008; Balkenhol and Waits 2009; Clark et al. 2010; Brehme et al. 2013). Direct mortality, or roadkill, is often associated with distance moved by organisms and their movement frequency and ability (Forman et al. 2003; Andrews et al. 2008). Research on some taxa has demonstrated dispersal limitations and subsequent decreased genetic relatedness due to clearings and roads (Stephens et al. 2013). Roads and habitat fragmentation, in general, create gene flow barriers for mammals, reptiles, and amphibians (Forman et al. 2003; Clark et al. 2010; Row et al.

2012; Stephens et al. 2013). Roads and trails, whether in-use or abandoned, reduce salamander abundance and fragment habitat (Semlitsch et al. 2007; Andrews et al. 2008). Environmental changes associated with roads and clearings may alter terrestrial plethodontid salamander abundance, distribution, genetic diversity, and movement activity (Kramer et al. 1993). Increased desiccation rates occur when the environment is changed and suitable moisture content is not maintained (Andrews et al. 2008). Roads negatively impact surrounding habitat and included organisms anywhere from 20-800m from the road edge, designated the “road-effect zone” (Forman and Alexander 1998; Marsh and Beckman 2004; Andrews et al. 2008; Semlitsch et al. 2012). Barriers and habitat edges can affect terrestrial salamander activity and movements, potentially causing genetic divergence between subpopulations (Gibbs 1998; Marsh and Beckman 2004; Marsh et al. 2008). In addition, competition for resources (e.g. food and habitat) and interactions between species may increase as habitat quality declines (Marsh et al. 2005; Kroschel et al. 2014).

Amphibians make excellent models for examining the effects of fragmentation, urbanization, and changing environmental conditions on ecosystems. Amphibians are particularly susceptible to habitat changes because they use cutaneous respiration and have limited movement ability, making them vulnerable to environmental changes and desiccation (Marsh and Beckman 2004; Marsh et al. 2005; Andrews et al. 2008). Such traits make amphibians appropriate habitat quality indicators (Welsh and Droege 2001; Marsh et al. 2005). Though previous studies have examined salamander abundance and condition in response to habitat fragmentation by roads (Marsh et al. 2004; Semlitsch et al. 2007), and others have assessed dispersal through fragmented areas by translocated individuals (Marsh et al. 2004; Marsh et al. 2005; Connette & Semlitsch 2013), no research to-date has examined foraging

movements in response to roads and trails. This project examined natural foraging patterns for individual salamanders near roads and trails. I analyzed movement patterns to examine the effects of roads and trails on salamander surface activity.

In this study, I assessed indirect impacts of roads and trails on movement behavior of two salamander species. Specifically, I examined movement data in four ways: 1) movement probability (move or no move), 2) total distance moved, 3) net distance moved, and 4) movement direction. Salamanders from the genus *Plethodon* are lungless, terrestrial, woodland salamanders that generally require cool, moist environments to survive (Connette and Semlitsch 2013). In this study, I focused on two plethodontid species: 1) the Peaks of Otter Salamander (*Plethodon hubrichti*, POS), a habitat specialist endemic to high elevations (>450m) along a 19km stretch of ridge bordering two counties in central Virginia (Kramer et al. 1993; Mitchell et al. 1996; Reichenbach and Sattler 2007; Kniewski and Reichenbach 2009); and 2) the Eastern Red-backed Salamander (*Plethodon cinereus*, RBS), a habitat generalist whose range extends from parts of North Carolina, north into Canada, and west as far as Minnesota, inhabiting forests at all elevations (Petranka 1998; Marsh et al. 2004; Marsh et al. 2008; Beane et al. 2010). The Peaks of Otter Salamander is globally imperiled, is listed as “vulnerable” on the IUCN Red List, is on the federal sensitive species list, and is under a US Fish and Wildlife Service candidate conservation agreement with the US Forest Service and Blue Ridge Parkway (Hammerson and Mitchell 2004; ECOS). I examined movement behavior for POS in allopatric sites and where it was sympatric with RBS to determine whether species interactions affected a salamander’s response to decreased habitat quality and fragmentation. I assessed movement behavior in response to salamander proximity to roads and trails for both species to determine whether a habitat specialist (POS) differed in movement behavior from a habitat generalist (RBS). I expected that

individuals in close proximity to roads and trails would move less than individuals at control sites. I also expected individuals to avoid treatment edges when foraging, due to decreased habitat quality, increased risk exposure to predation, and decreased canopy cover that would result in lower moisture and higher temperature. I expected individuals to move farther distances under cooler and moist conditions. Finally, I expected POS in sympatry with RBS to move less than POS in allopatric sites and that RBS would move more on average than POS. Studies of movement patterns help ecologists and wildlife managers better understand species life histories and environmental effects on activity. Additional studies on movement behavior will demonstrate long-term implications of habitat fragmentation.

METHODS

To examine the effects of roads and trails as barriers to terrestrial salamander movement, I used a natural experimental approach to quantify effects of habitat alteration and anthropogenic disturbance by gravel roads, paved roads, and hiking trails. I assumed impacts on salamander movements would decrease as distance from the barrier increased, indicating a linear effect (Marsh and Beckman 2004; Andrews et al. 2008). Further, I included candidate models that only contained conditional covariates known or suspected to drive plethodontid surface behavior, which allowed us to assess the validity of tracking methods. I assumed significant effects of conditional covariates (e.g., temperature, precipitation, and moon phase) would indicate that our tracking method did not adversely affect salamander movement behavior.

Study Sites

I assessed POS and RBS movement near roads and trails within the Jefferson National Forest along the Blue Ridge Parkway in Bedford and Botetourt Counties, Virginia. I sampled 16

sites, all located within sympatric and allopatric zones of POS distribution. Sites consisted of three treatments: 1) paved roads (i.e., Blue Ridge Parkway), 2) gravel roads (i.e. Forest Service roads), and 3) recreational trails (i.e., the Appalachian Trail). The edge effect is thought to influence terrestrial salamanders 80-100m from road edges (Marsh and Beckman 2004; Semlitsch et al. 2007). To ensure independence, control sites were located at least 200m from treatment edges and study sites were located approximately 200m apart.

Data collection

To locate salamanders during diurnal searches, I flipped natural cover objects in treatment (along roads and trails) and control sites between 5:00pm and sunset. I used fluorescent pigment powder (DayGlo ZQ-11 Aurora Pink; Figure 1) to track salamander movements. Fluorescent pigment powder has been used to track an array of organisms, including small mammals, reptiles, insects, and several amphibian species with no negative effects (Rittenhouse et al. 2006; Roe and Grayson 2008; Orlofske et al. 2009; Furman et al. 2011; Popescu and Hunter 2011). Once I captured a salamander, I measured mass (g) with a Ziploc bag clipped to a 10g Pesola spring scale, snout-vent length (SVL; mm) using a simplified “mander-masher” (Wise and Buchanan 1992) and ruler, determined sex by examining morphology (males exhibit square nose and swollen cloaca), dipped the posterior third of the salamander body in fluorescent pigment powder, returned them to their original capture location, and measured the salamander’s distance (m) from the road or trail using a measuring tape stretched from the salamander’s location to the closest point on the treatment edge. I returned to the site approximately six hours later to 1) determine if the salamander left its cover object (binary data; 1= moved; 0= did not move), 2) trace the pigment powder trail, 3) determine salamander trail direction (i.e. mark North and direction to treatment edge), and 4) photograph the fluorescent

powder trail. I used stake flags to mark initial cover objects for easy relocation when returning to trace the path. I recorded the capture location of each salamander with a handheld Global Positioning System (GPS; Trimble Juno SB; 5m accuracy). I recorded sampling covariates, including air temperature (°C) and percent relative humidity (Traceable temperature/humidity pen), and soil temperature (°C; Taylor 9878E thermometer probe) during each survey. Starting in mid-May 2014, I collected and tracked salamanders at two treatments per sampling night with 4-5 sampling nights per week. I began tracking individual movement trails around midnight, at the end of salamander peak foraging times. I avoided sampling during moderate to heavy rain, as the fluorescent powder would wash away, though I still attempted to sample on nights with forecasted rain. Sampling was conducted through June, or until no salamanders were detected at a site after 30 minutes of searching. In July, I recorded site covariates including canopy cover and elevation.



Figure 1: Fluorescent pigment powder applied to the belly and tail of a salamander as it looks in natural light (**A**) and under ultraviolet light (**B**).

I tracked individual salamander movements using a handheld ultraviolet flashlight (Northwest Marine; 405nm; 82mw) and traced each path using reflective rope. I placed a 1m²

PVC grid over the traced trail. If the trail was not contained within 1m², I added additional PVC pieces to create adjacent 1m squares. I used a small dry-erase board to note the individual's identification number and drew an arrow pointing toward the treatment edge (road or trail), where applicable. I used a handheld digital camera (Nikon D60 with 18-55mm DX VR lens or Canon PowerShot A2200) to photograph the 1-m² plot with all four corners visible (Figure 2). If the trail extended over multiple grids, I individually photographed each grid to minimize distortion. Photographs were oriented to magnetic north using an arrow placed on the ground, aligned with a compass. An additional arrow indicated the direction of the treatment.




Figure 2: Georectified image of a marked salamander movement trail using reflective rope with 1m² PVC grid, north arrow (orange), and dry-erase board with identification information and direction of treatment edge (blue arrow).

Data Analysis

I imported the digital photographs into ArcGIS 10.2, georectified each image, and transformed them to ensure edges of the PVC grid measured 1m. I opened the georectified images with ImageJ 1.48v and set the measurement scale to 1m using one of the horizontal edges

of the PVC grid from the image. I traced each salamander movement trail using the freehand draw tool and calculated the total distance traveled. I also used the line segment tool to measure net distance traveled from original cover object to the end of the trail.

I assessed movement direction for each salamander to determine if individuals near treatment edges were avoiding edges. Using ArcGIS 10.2, I measured each net movement angle (bearing) and compared the angle to the direction toward each treatment edge (randomized angles for control sites). I then divided the data into three equal directions: toward, away, or parallel (Figure 3).




Figure 3: Diagram of movement direction split into three equal bearings for analysis.

Candidate Models. I examined the effects of abiotic conditions, moon phase, treatment, and sympatry on salamander movement (Table 1). Prior to analyzing individual models, I used correlation analysis to assess collinearity of predictors, removing correlated variables ($r \geq 0.70$) from the analysis. Natural drivers included soil temperature, relative humidity, percent cloud cover, precipitation totals for the sampling day plus one day prior, as well as precipitation totals for the sampling day plus the three days prior. I measured soil temperature, relative humidity,

and percent cloud cover at the time of sampling, and precipitation totals were gathered from the National Oceanic and Atmospheric Administration's (NOAA) Global Historical Climatology Network (GHCN). The closest station to my study sites, and the station used for precipitation data, was Holcomb Rock (12-19km; GHCND:USC00444039). Soil temperature, humidity, and precipitation totals were z-standardized for analysis. Moon phase models included percent moon luminosity and moon altitude in relation to the horizon at 9:00pm, gathered from www.dailymoonposition.com. Treatment models included a categorical variable for the three treatments (trail, gravel, paved) and controls, as well as the distance from the original capture location to the treatment edge (randomized for control sites). Sympatry models included another categorical variable for whether the site was located in the region of POS allopatry or POS and RBS sympatry.

Table 1: Definition of predictor variables used in Peaks of Otter Salamander and Eastern Red-backed Salamander movement behavior models.

Variable	Definition
Cloud	Percent cloud cover at time of initial sampling (0.00-1.00)
Moon	Percent moon luminosity at 9:00pm (0.00-1.00)
Altitude	The degrees the moon is located above or below the horizon at 9:00pm (-90-90)
Precip2	Total precipitation for the two days prior to sampling, including the day of sampling, z-transformed
Precip4	Total precipitation for the four days prior to sampling, including the day of sampling, z-transformed
ST	Average soil temperature during initial sampling, z-transformed
RH	Average relative humidity during initial sampling, z-transformed
Sympatry	Categorical variable to identify site location (Allopatric or Sympatric)
Treatment	Categorical variable to identify site type (Paved, Gravel, Trail, or Control)
Edge	Distance (m) the salamander was found from the treatment edge, randomized for controls

Candidate models assessed natural and anthropogenic variables as sole predictors and in additive models (Table 4). I also included distance to treatment in a nested model to examine linear effects of treatment. I only used the sympatry model when comparing POS movement behavior between allopatric and sympatric sites to examine effects of potential competition.

Statistical Analysis. I conducted statistical analyses in SAS 9.4 to examine predictors of

movement. I examined fourteen models using anthropogenic and abiotic variables and ranked them using AICc model selection. I used models with ΔAICc less than 2.00 for inference. I assessed model fit by examining model deviance (Pearson chi-square/DF) for the most parameterized model. I used a generalized linear mixed model (GLIMMIX Procedure) in SAS 9.4, fit with a Laplace approximation (Waldron et al. 2013) to examine models using binary movement data (movement probability), continuous trail length data (total distance moved), net movement (straight-line distance from start to end) and movement direction (in relation to the treatment edge) as response variables in separate analyses. In all analyses, I specified site as a random effect to account for a lack of independence among observations from the same site. I used the response variables as the fixed effects for each model (Table 4). I also examined the effect of site location (allopatry or sympatry) on movement to determine if competition affected distance moved.

I used t-tests to examine interspecific differences in mass, SVL, and movement. I used logistic regression to examine binary movement data, modelling the probability a salamander moved from its initial cover object. I used Poisson regression to examine models with total distance or net distance as the response variable. For bearing analysis, which included a categorical response variable, I used a multinomial distribution and multinomial logistic regression. All candidate models were run to assess movement probability, total distance, net distance, and movement direction in two separate analyses: for the Peaks of Otter Salamander ($N = 141$), and for the Eastern Red-backed Salamander ($N = 46$).

RESULTS

I captured and marked 210 adult salamanders ($>30\text{mm}$; Kniewski and Reichenbach 2009)

in May and June, 2014. Heavy rain prevented tracking 23 individuals; therefore data collected from these individuals were not used in analyses. Of the 187 remaining individuals, 46 were RBS in sympatry with POS, 59 were sympatric POS, and 82 were allopatric POS. I recorded 118 individuals that moved from their original cover object (rock or log); the remaining 69 salamanders did not move from cover objects. The average mass and SVL of POS was 1.32g and 46.2mm, respectively. The average mass and SVL of RBS was 1.07g and 43.2mm, respectively (Table 2). Peaks of Otter salamanders were significantly larger than RBS in both mass ($t = 3.81$; $P = 0.0002$) and SVL ($t = 3.49$; $P = 0.0006$).

The average total distance moved by POS was 1.41 meters and the average distance moved by RBS was 1.04 meters (Figure 4; Table 3). Neither total distance moved ($t = 1.76$; $P = 0.081$) nor net distance moved ($t = 1.10$; $P = .2741$) were significantly different between species.

Table 2: Average mass (g) and snout-vent length (SVL; mm) for Peaks of Otter salamander (POS) and Eastern Red-backed salamander (RBS) with standard deviation and range of values.

Species	Mass (g)		SVL (mm)	
	Mean + SD	Range	Mean + SD	Range
POS	1.32 ± 0.41	0.40 - 2.50	46.2 ± 5.2	35.0 - 58.0
RBS	1.07 ± 0.31	0.60 - 1.80	43.2 ± 4.5	35.0 - 52.0




Table 3: Average total distance moved (m) and net distance moved (m) for Peaks of Otter salamanders (POS) and Eastern Red-backed salamanders (RBS), including range and standard deviation.

Species	Total Distance (m)		Net Distance (m)	
	Mean + SD	Range	Mean + SD	Range
POS	1.41 ± 1.07	0.10 - 6.50	0.86 ± 0.57	0.11 - 2.58
RBS	1.04 ± 0.84	0.18 - 3.49	0.73 ± 0.70	0.08 - 2.63

Correlation Analysis

Soil temperature was correlated with daily high air temperature ($r = 0.7386$; $P = <0.0001$), daily low air temperature ($r = 0.7877$; $P = <0.0001$), average daily air temperature ($r = 0.8577$; $P = <0.0001$), average three-day air temperature ($r = 0.7332$; $P = <0.0001$), and air temperature at sampling time ($r = 0.7419$; $P = <0.0001$). I retained soil temperature for analysis. Daily precipitation totals were correlated with two day precipitation totals ($r = 0.9727$; $P =$

<0.0001), thus I retained the two day precipitation totals for analysis. Finally, moon luminosity was correlated with moon altitude at midnight ($r = 0.9340$; $P = <0.0001$), but not moon altitude at 9:00pm, so I retained moon luminosity and moon altitude at 9:00pm for analysis. Removing correlated variables resulted in 11 variables retained for analyses, combined into 14 total candidate models (Table 4).

Table 4: Predictor variables used in candidate models to analyze Peaks of Otter Salamander and Eastern Red-backed Salamander movement behavior.

Model
Cloud
Moon
Moon + Altitude
Moon + Cloud
Moon + Cloud + Altitude
Precip2
Precip4
ST
ST + RH
ST + RH + Precip2
ST + RH + Precip4
Sympatry
Treatment
Treatment + Edge (Treatment)

I used the most parameterized model to examine model fit for each analysis. Movement analysis had great fit (Pearson $\chi^2/DF = 1.00$), whereas analysis of net distance moved was underdispersed (Pearson $\chi^2/DF = 0.69$) and analysis of total distance moved was slightly overdispersed (Pearson $\chi^2/DF = 1.12$).

Movement Probability

Peaks of Otter Salamander. I retained three models for inference out of 14 candidate models ($N = 141$; Table 5). The most important predictors of POS movement probability were moon luminosity ($\omega_i = 0.46$), moon altitude ($\omega_i = 0.36$), and Precip2 ($\omega_i = 0.23$; Table 4). The top

ranking model of movement probability for POS included moon luminosity and moon altitude. I failed to detect an effect of moon luminosity on POS movement probability ($\beta = -0.0663 \pm 0.0495$; $P = 0.1827$). Moon altitude was positively associated with movement probability ($\beta = 0.0129 \pm 0.0058$; $P = 0.0293$). The model containing precipitation totals for the two days prior to sampling (Precip2) as the sole predictor was the second highest ranking model, but I failed to detect a significant effect of Precip2 on POS movement probability ($\beta = 0.2862 \pm 0.1839$; $P = 0.1221$). The third supported model included moon luminosity, altitude, and cloud cover. I failed to detect significant effects of moon luminosity ($\beta = -0.0624 \pm 0.05286$; $P = 0.2200$), cloud cover ($\beta = 0.1904 \pm 0.5286$; $P = 0.7194$) or altitude ($\beta = 0.0121 \pm 0.0062$), though altitude approached significance ($P = 0.0530$).

Table 5: Candidate models examining Peaks of Otter Salamander (N = 141) movement probability (binary logistic regression), ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. ΔAICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta\text{AICc} < 2.00$ (in bold) were retained for inference.

Model	AICc	Δ AICc	ω_i
Moon + Altitude	189.95	0.00	0.26
Precip2	190.74	0.79	0.18
Moon + Cloud + Altitude	191.94	1.99	0.10
Cloud	192.12	2.17	0.09
Sympatry	192.59	2.64	0.07
Moon	192.94	2.99	0.06
ST + RH + Precip2	193.22	3.27	0.05
ST	193.45	3.50	0.05
Precip4	193.49	3.54	0.05
Moon + Cloud	193.79	3.84	0.04
ST + RH	193.89	3.94	0.04
ST + RH + Precip4	195.87	5.92	0.01
Treatment	197.33	7.38	0.01
Treatment + Edge(Treatment)	205.59	15.64	0.00

Red-backed Salamander. Analyses examining RBS movement probability resulted in four supported models out of 13 candidate models ($N = 46$; Table 6). Cloud cover was the top predictor of movement for RBS, accounting for 57% of model weights, followed by moon luminosity with 49% and soil temperature with 29% (Table 5). In the top ranking model, cloud cover and moon luminosity were negatively associated with RBS movement probability, though I failed to detect a significant effect of moon luminosity ($\beta = -2.0773 \pm 0.9887$; $P = 0.0427$; $\beta = -0.3128 \pm 0.408$; $P = 0.4484$, respectively). The model including cloud cover as the sole predictor of movement probability also indicated that movement probability increased as cloud cover decreased ($\beta = -2.0295 \pm 0.9573$; $P = 0.0408$). The third supported model included soil temperature and relative humidity as predictors of RBS movement probability. Relative humidity was positively associated with movement probability ($\beta = 10.1309 \pm 4.8078$; $P = 0.0421$), but I failed to detect a significant effect of soil temperature on RBS movement probability ($\beta = -0.1375 \pm 0.4520$; $P = 0.7626$). The fourth supported model included moon luminosity, moon altitude, and cloud cover as predictors of RBS movement probability. I failed to detect

significant effects of moon luminosity ($\beta = -0.3375 \pm 0.4219$; $P = 0.4290$) or moon altitude ($\beta = 0.0082 \pm 0.0119$; $P = 0.4943$) on RBS movement probability. Cloud cover was negatively associated with RBS movement probability, although the relationship approached significance ($\beta = -1.9301 \pm 1.0101$; $P = 0.0643$).

Table 6: Candidate models examining Eastern Red-backed Salamander (N = 46) movement probability (binary logistic regression), ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. ΔAICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta\text{AICc} < 2.00$ (in bold) were retained for inference.

Model	AICc	ΔAICc	ω_i
Moon + Cloud	53.77	0.00	0.26
Cloud	54.22	0.45	0.21
ST + RH	54.83	1.06	0.15
Moon + Cloud + Altitude	55.68	1.91	0.10
Moon	56.21	2.44	0.08
ST + RH + Precip2	56.57	2.80	0.06
Moon + Altitude	57.17	3.40	0.05
ST + RH + Precip4	57.33	3.56	0.04
ST	58.17	4.40	0.03
Precip2	60.84	7.07	0.01
Precip4	61.33	7.56	0.01
Treatment	63.01	9.24	0.00
Treatment + Edge(Treatment)	70.27	16.5	0.00

Total Distance Moved

Peaks of Otter Salamander. I retained four out of 14 candidate models examining total distance moved by POS (N = 141; Table 7). Cloud cover was the most important predictor of total distance moved by POS, accounting for 64% of model weights, followed by moon luminosity ($\omega_i = 0.58$) and moon altitude ($\omega_i = 0.38$; Table 7). The top-ranked model indicated cloud cover was positively associated with total distance moved ($\beta = 0.6281 \pm 0.2597$; $P =$

0.0171). The second supported model included cloud cover, moon altitude and moon luminosity as predictors of POS distance moved. I failed to detect significant effects of cloud cover ($\beta = 0.4414 \pm 0.2747$; $P = 0.1107$), moon altitude ($\beta = 0.0053 \pm 0.0032$; $P = 0.1039$), or moon luminosity ($\beta = -0.0546 \pm 0.0419$; $P = 0.1953$). The third supported model included moon luminosity and moon altitude as predictors of POS movement distance. I failed to detect a significant effect of moon luminosity on total distance moved ($\beta = -0.0646 \pm 0.0408$; $P = 0.1158$), but moon altitude was positively associated with total distance moved ($\beta = 0.0069 \pm 0.0030$; $P = 0.0243$). In the fourth supported model, total distance moved by POS was positively associated with cloud cover ($\beta = 0.5888 \pm 0.2604$; $P = 0.0255$), but I failed to detect a significant effect of moon luminosity ($\beta = -0.0404 \pm 0.0408$; $P = 0.3248$) on total distance moved.

Table 7: Candidate models examining Peaks of Otter Salamander (N = 141) total distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. ΔAICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta\text{AICc} < 2.00$ (in bold) were retained for inference.

Model	AICc	ΔAICc	ω_i
Cloud	358.16	0.00	0.27
Moon + Cloud + Altitude	358.59	0.43	0.21
Moon + Altitude	359.05	0.89	0.17
Moon + Cloud	359.14	0.98	0.16
ST + RH	361.50	3.34	0.05
Moon	362.21	4.05	0.04
ST + RH + Precip4	363.33	5.17	0.02
ST + RH + Precip2	363.57	5.41	0.02
ST	363.62	5.46	0.02
Sympatry	364.02	5.86	0.01
Precip2	364.05	5.89	0.01
Precip4	364.05	5.89	0.01
Treatment	366.32	8.16	0.00
Treatment + Edge(Treatment)	368.48	10.32	0.00

Red-backed Salamander. Two models out of 13 candidate models were retained for inference in the analysis examining total distance moved for RBS (N = 46; Table 8). Top predictors of total distance moved by all RBS included moon luminosity ($\omega_i = 0.50$), moon altitude ($\omega_i = 0.37$), and soil temperature ($\omega_i = 0.31$; Table 8). The top predictive model indicated moon altitude had a significant positive effect on total distance moved by RBS ($\beta = 0.0105 \pm 0.0049$; $P = 0.0391$). I failed to detect a significant effect of moon luminosity ($\beta = -0.1390 \pm 0.1743$; $P = 0.4304$) or soil temperature ($\beta = 0.2523 \pm 0.1946$; $P = 0.2028$) on total distance moved by RBS.

Table 8: Candidate models examining Eastern Red-backed Salamander (N = 46) total distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. ΔAICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta\text{AICc} < 2.00$ (in bold) were retained for inference.

Model	AICc	ΔAICc	ω_i
Moon + Altitude	103.36	0.00	0.29
ST	105.29	1.93	0.11
ST + RH	105.45	2.09	0.10
Moon	105.70	2.34	0.09
Moon + Cloud + Altitude	105.74	2.38	0.09
Cloud	106.19	2.83	0.07
Precip2	106.19	2.83	0.07
ST + RH + Precip4	106.30	2.94	0.07
Precip4	106.93	3.57	0.05
Moon + Cloud	107.38	4.02	0.04
ST + RH + Precip2	107.79	4.43	0.03
Treatment	110.77	7.41	0.01
Treatment + Edge(Treatment)	119.21	15.85	0.00

Net Distance Moved

Peaks of Otter Salamander. I retained four models for inference out of 14 models examining net distance moved for POS (N = 141; Table 9). Top predictors of net distance moved by POS included moon luminosity, cloud cover, and whether the individual was found in allopatry or sympatry, accounting for 54%, 22% and 16% of model weights, respectively (Table 8). I failed to detect a significant effect of any predictor on net distance moved among the supported models. I failed to detect a significant effect of moon luminosity on POS movement distance in any of the supported models (model 1: $\beta = 0.0459 \pm 0.0460$; $P = 0.3201$; model 3: $\beta = 0.0524 \pm 0.0470$; $P = 0.2669$; model 4: $\beta = 0.0434 \pm 0.0464$; $P = 0.3519$). I failed to detect an effect of sympatry on POS movement ($\beta = 0.1275 \pm 0.2468$; $P = 0.6063$). I failed to detect a significant effect of moon altitude or cloud cover, though both were positively associated with total distance moved by POS ($\beta = 0.0029 \pm 0.0037$; $P = 0.4379$; $\beta = 0.2322 \pm 0.3216$; $P = 0.4717$, respectively).

Table 9: Candidate models examining Peaks of Otter Salamander (N = 141) net distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. ΔAICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with $\Delta\text{AICc} < 2.00$ (in bold) were retained for inference.

Model	AICc	Δ AICc	ω_i
Moon	255.52	0.00	0.27
Sympatry	256.53	1.01	0.16
Moon + Altitude	257.01	1.49	0.13
Moon + Cloud	257.09	1.57	0.12
Cloud	258.15	2.63	0.07
Precip2	258.63	3.11	0.06
ST	258.80	3.28	0.05
Precip4	258.84	3.32	0.05
Treatment	260.66	5.14	0.02
ST + RH	260.91	5.39	0.02
Moon + Cloud + Altitude	261.04	5.52	0.02
ST + RH + Precip2	262.85	7.33	0.01
ST + RH + Precip4	263.05	7.53	0.01
Treatment + Edge(Treatment)	264.88	9.36	0.00

Red-backed Salamander. I retained three supported models out of 13 candidate models in the analysis of net distance moved by RBS (N = 46; Table 10). Moon luminosity was the top predictor of net distance moved by RBS, accounting for 49% of model weights, followed by moon altitude with 37% (Table 9). The top supported model for net distance moved by all RBS indicated a significant positive association with moon altitude ($\beta = 0.0120 \pm 0.0059$; $P = 0.0496$). I failed to detect a significant effect of moon luminosity on RBS movement ($\beta = -0.1475 \pm 0.2179$; $P = 0.5028$). I failed to detect a significant effect of soil temperature on net distance moved ($\beta = 0.2680 \pm 0.2318$; $P = 0.2551$). The third supported model included moon luminosity as a predictor of RBS movement. I failed to detect a significant effect of moon luminosity on RBS movement ($\beta = -0.1926 \pm 0.3459$; $P = 0.5811$).

Table 10: Candidate models examining Eastern Red-backed Salamander (N = 46) net distance moved (Poisson regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. Δ AICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with Δ AICc < 2.00 (in bold) were

retained for inference.

Model	AICc	Δ AICc	ω_i
Moon + Altitude	85.29	0.00	0.27
ST	86.87	1.58	0.12
Moon	87.22	1.93	0.10
Cloud	87.70	2.41	0.08
Moon + Cloud + Altitude	87.70	2.41	0.08
Precip2	87.96	2.67	0.07
ST + RH	88.00	2.71	0.07
Precip4	88.09	2.80	0.07
ST + RH + Precip4	88.72	3.43	0.05
Moon + Cloud	89.13	3.84	0.04
ST + RH + Precip2	90.22	4.93	0.02
Treatment	91.93	6.64	0.01
Treatment + Edge(Treatment)	100.94	15.65	0.00

Movement Direction

Peaks of Otter Salamander. One out of 14 candidate models was retained for inference in the analysis examining movement direction by POS (N = 85; Table 11). Precip4 was the top predictor of movement direction, accounting for 57% of model weights (Table 10). I failed to detect a significant effect of Precip4 on POS movement direction (toward versus away, $\beta = -0.4327 \pm 0.3645$; $P = 0.2408$; toward versus parallel, $\beta = 0.2811 \pm 0.2611$; $P = 0.2868$).

Table 11: Candidate models examining Peaks of Otter Salamander (N = 141) movement direction (multinomial logistic regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; sympatry = categorical variable for allopatric or sympatric site location; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. Δ AICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with Δ AICc < 2.00 (in bold) were retained for inference.

Model	AICc	Δ AICc	ω_i
Precip4	178.43	0.00	0.50
Precip2	180.51	2.08	0.18
ST + RH	181.94	3.51	0.09
ST + RH + Precip4	182.25	3.82	0.07
ST	183.51	5.08	0.04
ST + RH + Precip2	183.54	5.11	0.04
Cloud	184.18	5.75	0.03
Moon	184.43	6.00	0.02
Sympatry	186.40	7.97	0.01
Moon + Cloud	186.46	8.03	0.01
Moon + Altitude	186.97	8.54	0.01
Treatment	189.59	11.16	0.00
Moon + Cloud + Altitude	190.39	11.96	0.00
Treatment + Edge(Treatment)	205.33	26.90	0.00

Red-backed Salamander. I retained three models out of 13 candidate models for inference in the analysis examining RBS movement direction (N = 33; Table 12). Soil temperature was the most important predictor of RBS movement direction, accounting for 32% of model weights, followed by Precip2 ($\omega_i = 0.30$) and moon luminosity ($\omega_i = 0.21$; Table 11). I failed to detect a significant effect of Precip2 on RBS movement direction (toward versus away, $\beta = -3.5983 \pm 80.115$; $P = 0.9648$; toward versus parallel, $\beta = 13.156 \pm 37.309$; $P = 0.7299$). I failed to detect significant effects of other predictors, though supported models included soil temperature (toward versus away, $\beta = -0.3522 \pm 1.0082$; $P = 0.7317$; toward versus parallel, $\beta = 1.0235 \pm 0.6958$; $P = 0.1620$) and moon luminosity (toward versus away, $\beta = -0.3428 \pm 2.0767$; $P = 0.811$; toward versus parallel, $\beta = 2.3691 \pm 1.6416$; $P = 0.1695$).

Table 12: Candidate models examining Eastern Red-backed Salamander (N = 46) movement direction (multinomial logistic regression) ranked according to AICc model selection. Cloud = percent cloud cover; Moon = percent moon luminosity; Altitude = moon altitude; Precip2 = precipitation totals for two days prior to sampling; Precip4 = precipitation totals for four days prior to sampling; ST = soil temperature; RH = relative humidity; treatment = categorical variable for treatment type: paved, gravel, trail, or control; Edge = salamander distance from the treatment edge. Δ AICc = the difference between the AICc value for the current model and the lowest AICc score. ω_i = model weight. Models are listed in order of support. Models with Δ AICc

< 2.00 (in bold) were retained for inference.

Model	AICc	Δ AICc	ω_i
Precip2	71.70	0.00	0.30
ST	71.73	0.03	0.29
Moon	72.71	1.01	0.18
Cloud	73.89	2.19	0.10
Precip4	74.54	2.84	0.07
ST + RH	76.85	5.15	0.02
Moon + Cloud	77.43	5.73	0.02
Moon + Altitude	78.30	6.60	0.01
Treatment	80.24	8.54	0.00
ST + RH + Precip2	81.40	9.70	0.00
Moon + Cloud + Altitude	83.04	11.34	0.00
ST + RH + Precip4	83.22	11.52	0.00
Treatment + Edge(Treatment)	107.62	35.92	0.00

DISCUSSION

Many factors influence *Plethodon* salamander movement activity during peak foraging times. I failed to detect an effect of treatment or distance from treatment edge on movement behavior, indicating that the barriers I examined may not have an effect at the scale at which I examined them. Behavior was strongly influenced by natural predictors (e.g. cloud cover, relative humidity, and moon altitude), indicating decreased habitat quality near barriers may not drive salamander foraging behavior. Salamanders living in close proximity to barrier edges may have adapted to living in such conditions and therefore did not show changes in foraging behavior as compared to salamanders farther from a treatment edge. In addition, because salamander abundance is generally lower near road edges (Marsh and Beckman 2004; Semlitsch et al. 2007), individuals that live in these areas would most likely have lower competition with other individuals and potentially higher food and cover availability. Salamanders near road edges can be larger (Semlitsch et al. 2007), which may indicate individuals with higher body condition are able to live in these barrier edges. Future research should examine the time a salamander

spends on the surface and compare results at varying distances from barriers to determine habitat quality differences and road effects.

Though I failed to detect effects of treatment or distance from edge on salamander movements, my analyses indicated significant effects of abiotic predictors known to drive salamander surface behavior. Precipitation, soil temperature, cloud cover, and relative humidity were all top predictors of salamander movement, following conclusions from other studies (e.g., Taub 1961; Welsh and Droege 2001; Kniowski and Reichenbach 2009; MacNeil and Williams 2014). These results also indicate my use of fluorescent powder did not significantly affect movement behavior. Cloud cover and relative humidity were significant predictors of RBS movement probability. Salamanders were more likely to leave cover objects as relative humidity increased. My results also indicated RBS were more likely to move from cover objects when cloud cover decreased, which may suggest individuals moved underground or to more suitable cover objects. As cloud cover increased, however, total distance moved by POS increased, indicating salamanders might make longer foraging treks with increasing cloud cover.

Moon luminosity and moon altitude were important predictors of movement probability and movement distances for both salamander species. The model containing moon luminosity and moon altitude was consistently a top supported model. These relationships likely reflect risk exposure to predation during times of high luminosity. Plethodontid salamanders are not generally surface active during daytime hours and therefore moon luminosity may play an important role in nocturnal foraging behavior. Ralph (1957) found that not only are RBS more active at night, but they exhibit a reduction in activity when the moon's zenith (highest angle for a given day) occurs during peak times of activity. My results indicated that both species were more likely to leave cover objects and move longer distances when the moon was higher in the

sky, contradicting Ralph's results. Although not significant, moon luminosity (how much light the moon reflected) was consistently supported in top models, reinforcing my supposition that the moon is an important contributor of nocturnal salamander behavior. Salamander activity increased as the moon rose, as long as the moon reflected little light. Amphibian responses to lunar cycles are highly species-specific, with some exhibiting a negative response to moonlight and others exhibiting a positive response (Grant et al. 2013). Salamanders generally show a negative response to light. Wise and Buchanan (2006) found terrestrial salamanders might be photonegative as a response to desiccation (from increased temperature and decreased moisture levels) and predation risk. However, more light helps RBS find and orient toward prey, assisting their foraging ability (Wise and Buchanan 2006; Perry et al. 2008). In addition, salamanders in normal night light can exhibit photopositive responses and increased light can aid foraging (Wise and Buchanan 2006). One explanation for increased movement response to lunar position involves salamander diets. Both POS and RBS feed primarily on insects (Grant et al. 2013), which are more active at night with increased moonlight, supporting my observations of increased movement activity as the moon rose above the horizon. However, predation risk may reduce salamander movement with intense moonlight. Grant et al. (2013) found evidence that not all amphibians avoid activity at full moon despite higher predation risk. Amphibian predator avoidance may depend on visual acuity (Grant et al. 2013), resulting in a combination of factors that influence salamander risk assessment. Such factors might include visual acuity, prey availability, predator presence, and light intensity. Roberts and Liebgold (2008) found that terrestrial animals respond less to predators than aquatic animals. One means RBS and POS may reduce predation risk, even on moonlit nights, is by climbing vegetation (Roberts and Liebgold 2008). Plant-climbing activity is assumed to be a predator avoidance strategy, rather than a

foraging strategy, because the forest floor actually contains more prey for terrestrial salamanders (Roberts and Liebgold 2008). During nocturnal tracking, I observed POS and RBS perched on vegetation and climbing tree trunks up to waist high, usually on rainy or very wet nights.

I failed to detect a significant effect of treatment on individual movement direction, though out of 187 observations used for analyses, only one individual crossed a treatment edge (see Salamander Crossing note in Appendix A). I failed to detect an effect of proximity to treatment on the distance or direction an individual will move when foraging. However, on average, I observed RBS and POS move 1.31m over a 6-8 hour period, and I found individuals an average of 7.10m from treatment edges. Thus, observing foraging behavior over this short time frame might not be accurately demonstrating barrier edge effects. Although previous research has indicated salamander abundance declines near barrier edges (Marsh and Beckman 2004), I postulate that individuals in close proximity to roads and trails have habituated to environmental conditions and may not exhibit differences in foraging behavior as compared to individuals in more suitable, less fragmented, habitat.

Future studies should analyze salamander body condition and movement over longer time periods and through multiple seasons to better examine barrier avoidance. In addition, lunar effects on amphibians and specifically terrestrial salamanders has been poorly studied. Future research should continue to examine lunar position and luminosity as factors influencing terrestrial salamander movement behavior. Because my results indicated moonlight affects terrestrial salamander foraging behavior, light pollution from nearby homes and cities may alter salamander movement behavior, acting as an anthropogenic moon (Grant et al. 2013). In addition to these factors, future studies should analyze total time spent on the surface at varying distances from barriers to determine effects on behavior. The results of my study further support the use of

fluorescent pigment powder to quantify terrestrial plethodontid movement behavior. In addition, research examining the effects of anthropogenic features should include natural covariates to assess method validity and contribute insight into movement ecology.

LITERATURE CITED

- Andrews, K. M., J. W. Gibbons, and D. M. Jochimsen (2008). Ecological Effects of Roads on Amphibians and Reptiles: A Literature Review. *Herpetological Conservation* 3: 121-143.
- Balkenhol, N. and L. P. Waits (2009). Molecular road ecology: exploring the potential of genetics for investigating transportation impacts on wildlife. *Molecular Ecology* 18: 4151-4164.
- Beane, J. C.; A. L. Braswell, J. C. Mitchell, W. M. Palmer, and J. R. Harrison III (2010). *Amphibians and Reptiles of the Carolinas and Virginia, 2nd ed.* University of North Carolina Press, Chapel Hill, NC.
- Brehme, C. S., J. A. Tracey, L. R. McClanaghan, and R. N. Fisher (2013). Permeability of Roads to Movement of Scrubland Lizards and Small Mammals. *Conservation Biology* 27: 710-720.
- Clark, R. W., W. S. Brown, R. Stechert, and K. R. Zamudio (2010). Roads, Interrupted Dispersal, and Genetic Diversity in Timber Rattlesnakes. *Conservation Biology* 24:1059-1069.
- Connette, G. M. and R. D. Semlitsch (2013). Context-dependent movement behavior of woodland salamanders (*Plethodon*) in two habitat types. *Zoology* 116: 325-330.
- ECOS: Environmental Conservation Online System. *Peaks of Otter Salamander (Plethodon hubrichti)*. US Fish and Wildlife Service. ecos.fws.gov.
- Forman, R. T. and L. E. Alexander (1998). Roads and their major ecological effects. *Annual Review of Ecology and Systematics* 29: 207-231.
- Forman, R. T., D. Sperling, J. A. Bissonette, A. P. Clevenger, C. D. Cutshall, V. H. Dale, L. Fahrig, R. France, C. R. Goldman, K. Heanue, J. A. Jones, F. J. Swanson, T. Turrentine, and T.C. Winter (2003). *Road Ecology: Science and Solutions*. Island Press, Washington, D.C.
- Furman, B. L. S., B. R. Scheffers, and C. A. Paszkowski (2011). The Use of Fluorescent Powdered Pigments as a Tracking Technique for Snakes. *Herpetological Conservation and Biology* 6:473-478.
- Gibbs, J.P. (1998). Genetic structure of redback salamander *Plethodon cinereus* populations in continuous and fragmented forests. *Biological Conservation* 86: 77-81.
- Grant, R., T. Halliday, and E. Chadwick (2013). Amphibians' response to the lunar synodic cycle—a review of current knowledge, recommendations, and implications for conservation. *Behavioral Ecology* 24: 53-62.

- Hammerson, G. and J. Mitchell (2004). *Plethodon hubrichti*. The IUCN Red List of Threatened Species. www.iucnredlist.org.
- Kniowski, A. and N. Reichenbach (2009). The Ecology of the Peaks of Otter Salamander (*Plethodon hubrichti*) in sympatry with the Eastern Red-backed Salamander (*Plethodon cinereus*). *Herpetological Conservation and Biology* 4: 285-294.
- Kramer, P., N. Reichenbach, M. Hayslett, and P. Sattler (1993). Population Dynamics and Conservation of the Peaks of Otter Salamander, *Plethodon hubrichti*. *Journal of Herpetology* 27: 431-435.
- Kroschel, W. A., W. B. Sutton, C. J. W. McClure, T. K. Pauley (2014). Decline of the Cheat Mountain Salamander over a 32-Year Period and the Potential Influence of Competition from Sympatric Species. *Journal of Herpetology* 48: 415-422.
- MacNeil, J. E. and R. N. Williams (2014). Effects of Timber Harvests and Silvicultural Edges on Terrestrial Salamanders. *PLoS ONE* 9(12): e114683.
- Marsh, D. M., and N. G. Beckman (2004). Effects of forest roads on the abundance and activity of terrestrial salamanders. *Ecological Applications* 14: 1882-1891.
- Marsh, D. M., K. A. Thakur, K. C. Bulka and L. B. Clarke (2004). Dispersal and Colonization through Open Fields by a Terrestrial, Woodland Salamander. *Ecology* 85: 3396-3405.
- Marsh, D. M., G. S. Milam, N. P. Gorham, and N. G. Beckman (2005). Forest Roads as Partial Barriers to Terrestrial Salamander Movement. *Conservation Biology* 19: 2004-2008.
- Marsh, D. M., R. B. Page, T. J. Hanlon, R. Corritone, E. C. Little, D. E. Seifert, and P. R. Cabe (2008). Effects of roads on patterns of genetic differentiation in red-backed salamanders, *Plethodon cinereus*. *Conservation Genetics* 9: 603-613.
- Mazerolle, M. J. (2001). Amphibian Activity, Movement Patterns, and Body Size in Fragmented Peat Bogs. *Journal of Herpetology* 35:13-20.
- Mitchell, J. C., J. A. Wicknick, and C. D. Anthony (1996). Effects of Timber Harvesting Practices on Peaks of Otter Salamander (*Plethodon hubrichti*) Populations. *Amphibian and Reptile Conservation* 1: 15-19.
- Orlofske, S. A., K. L. Grayson, and W. A. Hopkins (2009). The Effects of Fluorescent Tracking Powder on Oxygen Consumption in Salamanders Using Either Cutaneous or Bimodal Respiration. *Copeia* 2009: 623-627.
- Perry, G., B. W. Buchanan, R. N. Fisher, M. Salmon, and S. E. Wise (2008). Effects of Artificial Night Lighting on Amphibians and Reptiles in Urban Environments. *Herpetological Conservation* 3: 239-256.

- Petranka, J. W. (1998). *Salamanders of the United States and Canada*. Smithsonian Institutional Press, Washington, D.C.
- Popescu, V. D. and M. L. Hunter, Jr. (2011). Clear-cutting affects habitat connectivity for a forest amphibian by decreasing permeability to juvenile movements. *Ecological Adaptation* 21: 1283-1295.
- Ralph, C. L. (1957). A Diurnal Activity Rhythm in *Plethodon cinereus* and Its Modification by an Influence Having a Lunar Frequency. *Biological Bulletin* 113: 188-197.
- Reichenbach, N. and P. Sattler (2007). Effects of Timbering on *Plethodon hubrichti* over Twelve Years. *Journal of Herpetology* 41: 622-629.
- Rittenhouse, T. A. G., T. T. Altnether, and R. D. Semlitsch (2006). Fluorescent Powder Pigments as a Harmless Tracking Method for Ambystomatids and Ranids. *Herpetological Review* 37: 188-191.
- Roberts, A. M. and E. B. Liebgold (2008). The effects of perceived mortality risk on habitat selection in a terrestrial salamander. *Behavioral Ecology* 19: 621-626.
- Roe, A. W. and K. L. Grayson (2008). Terrestrial Movements and Habitat Use of Juvenile and Emigrating Adult Eastern Red-spotted Newts, *Notophthalmus viridescens*. *Journal of Herpetology* 42: 22-30.
- Row, J. R., G. Blouin-Demers, and S. C. Loughheed (2012). Movements and Habitat Use of Eastern Foxsnakes (*Pantherophis gloydi*) in Two Areas Varying in Size and Fragmentation. *Journal of Herpetology* 46: 94-99.
- Semlitsch, R. D., T. J. Ryan, K. Hamed, M. Chatfield, B. Drehman, N. Pekarek, M. Spath, and A. Watland (2007). Salamander Abundance along Road Edges and within Abandoned Logging Roads in Appalachian Forests. *Conservation Biology* 21: 159-167.
- Semlitsch, R. D. S. Ecrement, A. Fuller, K. Hammer, J. Howard, C. Krager, J. Mozeley, J. Ogle, N. Shipman, J. Speier, M. Walker, and B. Walters (2012). Natural and anthropogenic substrates affect movement behavior of the Southern Graycheek Salamander (*Plethodon metclafi*). *Canadian Journal of Zoology* 90: 1128-1135.
- Stephens, H. C., C. Schmuki, C. P. Burridge, and J. M. O'Reilly-Wapstra (2013). Habitat fragmentation in forests effects relatedness and spatial genetic structure of a native rodent, *Rattus lutreolus*. *Austral Ecology* 38: 568-580.
- Taub, F. (1961). The Distribution of the Red-backed Salamander, *Plethodon c. cinereus*, within the Soil. *Ecology* 42: 681-698.
- Waldron, J. L., S. M. Welch, S. H. Bennett, W. G. Kalinowsky, and T. A. Mousseau (2013). Life history constraints contribute to the vulnerability of a declining North American

- rattlesnake. *Biological Conservation* 159: 530-538.
- Welsh, H. H., Jr. and S. Droege (2001). A Case for Using Plethodontid Salamanders for Monitoring Biodiversity and Ecosystem Integrity of North American Forests. *Conservation Biology* 15: 558-569.
- Wise, S. E. and B. W. Buchanan (1992). An Efficient Method for Measuring Salamanders. *Herpetological Review* 23: 56-57.
- Wise, S. E., and B. W. Buchanan (2006). Influence of artificial illumination on the nocturnal behavior and physiology of salamanders. In C. Rich and T. Longcore (eds.), *Ecological Consequences of Artificial Night Lighting*. Island Press, Washington, D.C. Pp. 221-251.

APPENDIX A

Salamander Crossing Note

Out of 187 salamanders tracked during my study, 49 individuals were located along hiking trails. On June 9, 2014, I tracked a female Peaks of Otter Salamander (42mm SVL; 1.2g), in an allopatric zone, from its original cover rock, across the hiking trail, to the other side and underneath a downed tree. I originally found the individual 0.40m from the edge of the trail and it moved a total distance of 3.73m, for a net distance of 2.41m. At time of original capture, the air temperature was 19.6°C, soil temperature was 16.1°C, and the relative humidity was 72%. When I returned and tracked the salamander movement trail, the air temperature had dropped to 17.7°C, soil temperature was still at 16.0°C, and humidity had increased to 81%. I documented cloud cover at 70%. This was the only salamander during my study that crossed a barrier.

Salamander Microhabitat Use

During my research, I observed and documented microhabitat use for the Peaks of Otter Salamander and the Eastern Red-backed Salamander. I noted both species perched on ferns or small shrubs, approximately one foot above the ground (Figure 5). These observations generally occurred on rainy nights. In addition, I observed both species climbing tree trunks (Figure 6), sometimes pressed into the deep folds of oak tree bark, up to one meter above the ground. Again, these observations generally occurred on rainy nights. Finally, though some salamander movement trails ended under other cover objects (rocks and logs), most trails ended when the individual followed small holes in the ground, usually near the base of plants and trees. These notes may provide insight into microhabitat use by both species.



Figure 5: A Peaks of Otter Salamander perched on a small plant, covered in fluorescent pigment powder used for tracking.



Figure 6: A Peaks of Otter Salamander climbing the trunk of a tree.



Animal Resource Facility

DATE: July 16, 2014

TO: Jayme Waldron, PhD
FROM: Marshall University IACUC

IACUC #: MU
PROJECT TITLE: [581002-3] Effects of trails and roads on Peaks of Otter Salamander
(Plethodon hubrichti) movement behavior
SUBMISSION TYPE: New Project

ACTION: APPROVED
APPROVAL DATE: July 16, 2014
EXPIRATION DATE: December 31, 2014
REVIEW TYPE: Full Committee Designated Member Review

Thank you for your submission of Revision materials for this research project. The Marshall University IACUC has APPROVED your submission. All research must be conducted in accordance with this approved submission.

This submission has received Full Committee Review of 581002-1 and Designated Member Review of 581002-3.

Please note that any revision to previously approved materials must be approved by this committee prior to initiation. Please use the appropriate revision forms for this procedure.

Please report all NON-COMPLIANCE issues regarding this project to this committee.

This project requires Continuing Review by this office on an annual basis. Please use the appropriate renewal forms for this procedure.

If you have any questions, please contact Monica Valentovic at (304) 696-7332 or valentov@marshall.edu. Please include your project title and reference number in all correspondence with this committee.

Monica A. Valentovic, Ph.D.
Chairperson, IACUC



Animal Resource Facility

DATE: May 19, 2014

TO: Jayme Waldron, PhD
FROM: Marshall University IACUC

IACUC #: 580
PROJECT TITLE: [599250-2] Effects of trails and roads on Eastern Red-backed Salamander (Plethodon cinereus) movement behavior.
SUBMISSION TYPE: Revision

ACTION: APPROVED
APPROVAL DATE: May 19, 2014
EXPIRATION DATE: December 19, 2014
REVIEW TYPE: Designated Member Review

Thank you for your submission of Revision materials for this research project. The Marshall University IACUC has APPROVED your submission. All research must be conducted in accordance with this approved submission.

This submission has received Designated Member Review.

Please note that any revision to previously approved materials must be approved by this committee prior to initiation. Please use the appropriate revision forms for this procedure.

Please report all NON-COMPLIANCE issues regarding this project to this committee.

This project requires Continuing Review by this office on an annual basis. Please use the appropriate renewal forms for this procedure.

If you have any questions, please contact Monica Valentovic at (304) 696-7332 or valentov@marshall.edu. Please include your project title and reference number in all correspondence with this committee.

Monica A. Valentovic, Ph.D.
Chairperson, IACUC